

## PRODUCT DATA SHEET

### lab0212 Anti-Mouse IgG (H+L), Peroxydase conjugate antibody



#### Product specifications

<b>Immunogen</b>	IgG(H+L)		
<b>Form</b>	F(ab') <sub>2</sub> fragments	<b>Target</b>	mouse
<b>Raised in</b>	Rabbit	<b>Labelling</b>	HRP
<b>Conjugate properties</b>	Peroxidase labelling has the advantage of being sensitive and permanent. Peroxidase substrates are varied: precipitating substrates for western blot or immuno-histochemistry (DAB, AEC, 4CN, TMB), soluble substrates for ELISA (TMB, OPD, ABTS) and chemiluminescent substrates (luminol). However, peroxidase labelling has some disadvantages: Peroxidase is rendered inactivated by oxygen, hypochloric acid and sodium azide. It is therefore advisable to use highly purified water. The presence of endogenous peroxidase in certain tissues like macrophage, red blood cells, bone marrow cells and plants could increase the background.		
<b>Applications</b>	The F(ab') <sub>2</sub> is bivalent fragments of immunoglobulins, i.e. including two sites for the fixation of the antigen. They are prepared by enzymatic digestion (pepsin). Like our others antibodies, they are purified by affinity chromatography and are thus particularly specific. F(ab') <sub>2</sub> fragments have certain advantages with respect to the entire antibodies in flow cytometry, immuno-histochemistry and immuno-cytology. On the one hand their specificity is increased since they can no longer be bound in a non-specific way by their Fc part as entire Ig's are. On the other hand their use also increases the sensitivity of the technique. In fact, their small size enhances their penetration into tissues and cells as well as their availability for the antigens (particularly for Fab). Finally, conjugated Fab's have a label/antigenic ratio that is twice as high compared to antibodies (except fluorescent conjugates).		
<b>Working dilutions</b>	Optimal dilution should be determined by the end user The following are guidelines only : - ELISA : 1/500 to 1/5 000 - Immunoblot : 1/100 to 1/1 000 - Immunocytochemistry : 1/20 to 1/100 - Immunohistochemistry : 1/20 to 1/100		

#### Packaging specifications

<b>Packaging</b>	1 ml
<b>Appearance</b>	Liquid
<b>Constituents</b>	TBS, 50% Glycerol
<b>Preservative</b>	None
<b>Storage</b>	Short term storage : +4°C Long term storage : -20°C
<b>Recommendations</b>	Product in glycerol, do not freeze Do not distribute the HRP conjugated in plastics (e.g. eppendorf) so as to avoid a loss of activity

**Product type**  
Secondary antibody

**Clonality**  
Polyclonal Antibody

**Related product**  
opr0002 ; opr0008 ;  
opr0009

**Warning**  
This is a laboratory reagent. It is not to be administered to human or animals nor be used as a drug

updated 04/10/06

version B

**Covalab France**  
11, Avenue Albert Einstein  
69100 Villeurbanne - France  
Phone +33 (0) 437 654 230  
Fax +33 (0) 437 289 416  
Email pa@covalab.com  
www.covalab.com

**Covalab - UK Ltd**  
St John's Innovation Centre  
Unit 75, Cowley Road  
Cambridge, CB40WS - UK  
Phone +44 (0) 1223 421055  
Fax +44 (0) 1223 420844  
Email enquiries@covalab.co.uk  
www.covalab.co.uk